HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) safely and effectively. See full prescribing information for Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound).

PACLITAXEL Protein-Bound Particles for Injectable Suspension (Albumin-Bound), for intravenous use

Initial U.S. Approval: 2005

WARNING: SEVERE MYELOSUPPRESSION

See full prescribing information for complete boxed warning.

- **Do not administer Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy to patients with baseline neutrophil counts of less than 1,500 cells/mm³ (*see Contraindications (4)*).**
- **Monitor for neutropenia, which may be severe and result in infection or sepsis (*see Warnings and Precautions (5.1, 5.3)*).**
- **Perform frequent complete blood cell counts on all patients receiving Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) (*see Warnings and Precautions (5.1, 5.3)*).**

RECENT MAJOR CHANGES

Dosage and Administration (2.1, 2.7)	8/2020
Contraindications (4)	8/2020
Warnings and Precautions (5.1, 5.2)	8/2020

INDICATIONS AND USAGE

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is a microtubule inhibitor indicated for the treatment of:

- Metastatic breast cancer, after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless clinically contraindicated. (1.1)
- Locally advanced or metastatic non-small cell lung cancer (NSCLC), as first-line treatment in combination with carboplatin, in patients who are not candidates for curative surgery or radiation therapy. (1.2)
- Metastatic adenocarcinoma of the pancreas as first-line treatment, in combination with gemcitabine. (1.3)

DOSAGE AND ADMINISTRATION

- **Do not substitute Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) for other paclitaxel products. (2.1)**
- Extravasation: Closely monitor the infusion site for extravasation and infiltration. (2.1)
- Metastatic Breast Cancer (MBC): Recommended dosage of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is 260 mg/m² intravenously over 30 minutes every 3 weeks. (2.2)
- Non-Small Cell Lung Cancer (NSCLC): Recommended dosage of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is 100 mg/m² intravenously over 30 minutes on Days 1, 8, and 15 of each 21-day cycle; administer carboplatin on Day 1 of each 21-day cycle immediately after Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). (2.2)
- Adenocarcinoma of the Pancreas: Recommended dosage of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is 125 mg/m² intravenously over 30-40 minutes on Days 1, 8, and 15 of each 28-day cycle; administer gemcitabine on Days 1, 8, and 15 of each 28-day cycle immediately after Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). (2.4)
- Use in Patients with Hepatic Impairment: Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is not recommended for use in patients with AST > 10 x ULN; or bilirubin > 5 x ULN or with metastatic adenocarcinoma of the pancreas who have moderate to severe hepatic impairment. For MBC or NSCLC, reduce starting dose in patients with moderate to severe hepatic impairment. (2.5)

Revised: 8/2020

PACTXPI.012.RPI.012 8/2020
Revised: August 2020

112462

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)

Rx only

(Patient Information Enclosed)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: SEVERE MYELOSUPPRESSION

1 INDICATIONS AND USAGE
1.1 Metastatic Breast Cancer
1.2 Non-Small Cell Lung Cancer
1.3 Adenocarcinoma of the Pancreas

2 DOSAGE AND ADMINISTRATION
2.1 Important Administration Instructions
2.2 Recommended Dosage for Metastatic Breast Cancer
2.3 Recommended Dosage for Non-Small Cell Lung Cancer
2.4 Recommended Dosage for Adenocarcinoma of the Pancreas
2.5 Dosage Modifications for Hepatic Impairment
2.6 Dosage Modifications for Adverse Reactions
2.7 Preparation for Intravenous Administration
2.8 Stability

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS
5.1 Severe Myelosuppression
5.2 Severe Neuropathy
5.3 Sepsis
5.4 Pneumonitis
5.5 Severe Hypersensitivity
5.6 Use in Patients with Hepatic Impairment
5.7 Albumin (Human)
5.8 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Renal Impairment
8.7 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES
14.1 Metastatic Breast Cancer
14.2 Non-Small Cell Lung Cancer
14.3 Adenocarcinoma of the Pancreas

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)

WARNING: SEVERE MYELOSUPPRESSION

- **Do not administer Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy to patients with baseline neutrophil counts of less than 1,500 cells/mm³ (*see Contraindications (4)*).**
- **Monitor for neutropenia, which may be severe and result in infection or sepsis (*see Warnings and Precautions (5.1, 5.3)*).**
- **Perform frequent complete blood cell counts on all patients receiving Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) (*see Warnings and Precautions (5.1, 5.3)*).**

1 INDICATIONS AND USAGE

1.1 Metastatic Breast Cancer

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is indicated for the treatment of breast cancer after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless clinically contraindicated.

1.2 Non-Small Cell Lung Cancer

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is indicated for the first-line treatment of locally advanced or metastatic non-small cell lung cancer, in combination with carboplatin, in patients who are not candidates for curative surgery or radiation therapy.

1.3 Adenocarcinoma of the Pancreas

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is indicated for the first-line treatment of patients with metastatic adenocarcinoma of the pancreas, in combination with gemcitabine.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

DO NOT SUBSTITUTE FOR OR WITH OTHER PACLITAXEL FORMULATIONS. Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) has different dosage and administration instructions from other paclitaxel products.

Closely monitor the infusion site for extravasation or drug infiltration during administration. Limiting the infusion of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) to 30 minutes may reduce the risk of infusion-related reactions *[see Adverse Reactions (6.2)]*.

Consider premedication in patients who have had prior hypersensitivity reactions to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). Do not re-challenge patients who experience a severe hypersensitivity reaction to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Contraindications (4) and Warnings and Precautions (5.5)]*.

2.2 Recommended Dosage for Metastatic Breast Cancer

After failure of combination chemotherapy for metastatic breast cancer or relapse within 6 months of adjuvant chemotherapy, the recommended regimen for Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is 260 mg/m² administered intravenously over 30 minutes every 3 weeks.

2.3 Recommended Dosage for Non-Small Cell Lung Cancer

The recommended dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is 100 mg/m² administered as an intravenous infusion over 30 minutes on Days 1, 8, and 15 of each 21-day cycle. Administer carboplatin on Day 1 of each 21-day cycle immediately after Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Clinical Studies (14.2)]*.

2.4 Recommended Dosage for Adenocarcinoma of the Pancreas

The recommended dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is 125 mg/m² administered as an intravenous infusion over 30-40 minutes on Days 1, 8, and 15 of each 28-day cycle. Administer gemcitabine immediately after the Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) on Days 1, 8, and 15 of each 28-day cycle *[see Clinical Studies (14.3)]*.

2.5 Dosage Modifications for Hepatic Impairment

For patients with moderate or severe hepatic impairment, reduce the starting dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) as shown in Table 1.

Table 1: Recommendations for Starting Dose in Patients with Moderate and Severe Hepatic Impairment

	AST Levels		Bilirubin Levels	Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) Dose*		
				MBC	NSCLC†	Adenocarcinoma of Pancreas†
Moderate	< 10 x ULN	AND	> 1.5 to ≤ 3 x ULN	200 mg/m²	80 mg/m²	not recommended
Severe	< 10 x ULN	AND	> 3 to ≤ 5 x ULN	200 mg/m²	80 mg/m²	not recommended
	> 10 x ULN	OR	> 5 x ULN	not recommended	not recommended	not recommended

AST = Aspartate Aminotransferase; MBC = Metastatic Breast Cancer; NSCLC = Non-Small Cell Lung Cancer; ULN = Upper limit of normal.
* Dosage recommendations are for the first course of therapy. The need for further dose adjustments in subsequent courses should be based on individual tolerance.
† A dose increase to 260 mg/m² for patients with metastatic breast cancer or 100 mg/m² for patients with non-small cell lung cancer in subsequent courses should be considered if the patient tolerates the reduced dose for two cycles.

Patients with bilirubin levels above the upper limit of normal were excluded from clinical trials for pancreatic or lung cancer.

2.6 Dosage Modifications for Adverse Reactions

Metastatic Breast Cancer

Patients who experience severe neutropenia (neutrophils less than 500 cells/mm³ for a week or longer) or severe sensory neuropathy during Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy should have dose reduced to 220 mg/m² for subsequent courses of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). For treatment of severe neutropenia or severe sensory neuropathy, additional dose reduction should be made to 180 mg/m² (≥ 20%) in NSCLC and adenocarcinoma of the pancreas or neutropenia, fatigue, peripheral neuropathy, nausea, and stomatitis.

The most common (≥ 20%) adverse reactions of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in adenocarcinoma of the pancreas are neutropenia, fatigue, peripheral neuropathy, nausea, peripheral edema, diarrhea, pyrexia, vomiting, decreased appetite, rash, and dehydration. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact TWI Pharmaceuticals USA, Inc. at 1-844-518-2989 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Use caution when concomitantly administering Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) with inhibitors or inducers of CYP3A4. (7)

USE IN SPECIFIC POPULATIONS

- Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 8/2020

- Dose Reductions for Adverse Reactions: Dose reductions or discontinuation may be needed based on severe hematologic, neurologic, cutaneous, or gastrointestinal toxicities. (2.6)

DOSAGE FORMS AND STRENGTHS

For injectable suspension: white to yellow, sterile, lyophilized powder containing 100 mg of paclitaxel formulated as albumin-bound particles in single-dose vial for reconstitution. (3)

CONTRAINDICATIONS

- Neutrophil counts of <1,500 cells/mm³. (4)
- Severe hypersensitivity reactions to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). (4)

WARNINGS AND PRECAUTIONS

- Sensory neuropathy occurs frequently and may require dose reduction or treatment interruption. (5.2)
- Sepsis occurred in patients with or without neutropenia who received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine; interrupt Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine until sepsis resolves, and if neutropenia, until neutrophils are at least 1500 cells/mm³, then resume treatment at reduced dose levels. (5.3)
- Pneumonitis occurred with the use of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine; permanently discontinue treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine. (5.4)
- Severe hypersensitivity reactions with fatal outcome have been reported. Do not rechallenge with this drug. (4, 5.5)
- Exposure and toxicity of paclitaxel can be increased in patients with hepatic impairment; monitor patients with mild hepatic impairment. (2.5, 5.6)
- Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) contains albumin derived from human blood, which has a theoretical risk of viral transmission. (5.7)
- Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) can cause fetal harm. Advise patients of the potential risk to a fetus and to use effective contraception. (5.8, 8.1, 8.3)

ADVERSE REACTIONS

- The most common adverse reactions (≥ 20%) in metastatic breast cancer are alopecia, neutropenia, sensory neuropathy, abnormal ECG, fatigue/asthenia, myalgia/arthralgia, AST elevation, alkaline phosphatase elevation, anemia, nausea, infections, and diarrhea. (6.1)
- The most common adverse reactions (≥ 20%) in NSCLC are anemia, neutropenia, thrombocytopenia, alopecia, peripheral neuropathy, nausea, and fatigue. (6.1)
- The most common (≥ 20%) adverse reactions of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in adenocarcinoma of the pancreas are neutropenia, fatigue, peripheral neuropathy, nausea, peripheral edema, diarrhea, pyrexia, vomiting, decreased appetite, rash, and dehydration. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact TWI Pharmaceuticals USA, Inc. at 1-844-518-2989 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Use caution when concomitantly administering Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) with inhibitors or inducers of CYP3A4. (7)

USE IN SPECIFIC POPULATIONS

- Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 8/2020

3 DOSAGE FORMS AND STRENGTHS

For injectable suspension: white to yellow, sterile, lyophilized powder containing 100 mg of paclitaxel formulated as albumin-bound particles in single-dose vial for reconstitution.

4 CONTRAINDICATIONS

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is contraindicated in patients with:

- Baseline neutrophil counts of < 1,500 cells/mm³ *[see Warnings and Precautions (5.1)]*.
- A history of severe hypersensitivity reactions to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Warnings and Precautions (5.5)]*.

5 WARNINGS AND PRECAUTIONS

5.1 Severe Myelosuppression

Severe myelosuppression (primarily neutropenia) is dose-dependent and is the dose-limiting toxicity of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). In clinical studies, Grade 4 neutropenia occurred in 34% of patients with metastatic breast cancer (MBC), 47% of patients with non-small cell lung cancer (NSCLC), and 38% of patients with pancreatic cancer.

Monitor for severe neutropenia and thrombocytopenia by performing complete blood cell counts frequently, including prior to dosing on Day 1 (for MBC) and Days 1, 8, and 15 (for NSCLC and for pancreatic cancer). Do not administer Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) to patients with baseline absolute neutrophil counts (ANC) of less than 1,500 cells/mm³ *[see Contraindications (4)]*.

In the case of severe neutropenia (<500 cells/mm³ for seven days or more) during a course of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy, reduce the dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in subsequent courses in patients with either MBC or NSCLC.

In patients with MBC, resume treatment with every-3-week cycles of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) after ANC recovers to at least 1,500 cells/mm³ and platelets recover to a level >100,000 cells/mm³. In patients with NSCLC, resume treatment with every-3-week cycles at reduced doses for both weekly Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and every-3-week carboplatin after ANC recovers to at least 1500 cells/mm³ and platelet count of at least 100,000 cells/mm³ on Day 1 or to an ANC of at least 500 cells/mm³ and platelet count of at least 50,000 cells/mm³ on Days 8 or 15 of the cycle *[see Dosage and Administration (2.6)]*.

In patients with adenocarcinoma of the pancreas, withhold Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine if ANC is less than 1500 cells/mm³ or platelets are less than 50,000 cells/mm³ and delay initiation of the next cycle of therapy until ANC is at least 1500 cells/mm³ and platelet is less than 100,000 cells/mm³ on Day 1 of the cycle. Resume treatment with appropriate dose reduction if recommended *[see Dosage and Administration (2.6)]*.

5.2 Severe Neuropathy

Sensory neuropathy is dose- and schedule-dependent *[see Adverse Reactions (6.1)]*. If ≥ Grade 3 sensory neuropathy develops, withhold Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) treatment until resolution to Grade 1 or 2 for metastatic breast cancer or Grade 1 for NSCLC and pancreatic cancer followed by a dose reduction for all subsequent courses. Patients with NSCLC and pancreatic cancer followed by a dose reduction for all subsequent courses. Patients with NSCLC and pancreatic cancer followed by a dose reduction for Injectable Suspension (Albumin-Bound) *[see Dosage and Administration (2.6)]*.

5.3 Sepsis

Sepsis occurred in 5% of patients with or without neutropenia who received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine. Biliary obstruction or presence of biliary stent were risk factors for severe or fatal sepsis.

If a patient becomes febrile (regardless of ANC) initiate treatment with broad spectrum antibiotics. For febrile neutropenia, interrupt Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine if ANC is less than 500 cells/mm³ and ANC ≥ 1500, then resume treatment at reduced dose levels *[see Dosage and Administration (2.6)]*.

5.4 Pneumonitis

Pneumonitis, including some cases that were fatal, occurred in 4% of patients receiving Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine.

Monitor patients for signs and symptoms of pneumonitis and interrupt Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine during evaluation of suspected pneumonitis. After ruling out infectious etiology and upon making a diagnosis of pneumonitis, permanently discontinue treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine.

5.5 Severe Hypersensitivity

Severe and sometimes fatal hypersensitivity reactions, including anaphylactic reactions, have been reported. Do not rechallenge patients who experience a severe hypersensitivity reaction to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) with this drug *[see Contraindications (4)]*.

Cross-hypersensitivity between Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and other taxane products has been reported and may include severe reactions such as anaphylaxis. Closely monitor patients with a previous history of hypersensitivity to other taxanes during initiation of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy.

5.6 Use in Patients with Hepatic Impairment

The exposure and toxicity of paclitaxel can be increased in patients with hepatic impairment. Closely monitor patients with mild hepatic impairment for severe myelosuppression.

The most common adverse reactions (≥ 20%) with single-agent use of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) are hematologic toxicity, neuropathy, nausea, sensory neuropathy, abnormal ECG, fatigue/asthenia, myalgia/arthralgia, AST elevation, alkaline phosphatase elevation, anemia, nausea, infections, and diarrhea *[see Adverse Reactions (6.1)]*.

Monitor for severe neutropenia and thrombocytopenia by performing complete blood cell counts frequently, including prior to dosing on Day 1 (for MBC) and Days 1, 8, and 15 (for NSCLC and for pancreatic cancer). Do not administer Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) to patients with baseline absolute neutrophil counts (ANC) of less than 1,500 cells/mm³ *[see Contraindications (4)]*.

In the case of severe neutropenia (<500 cells/mm³ for seven days or more) during a course of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy, reduce the dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in subsequent courses in patients with either MBC or NSCLC.

In patients with MBC, resume treatment with every-3-week cycles of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) after ANC recovers to at least 1,500 cells/mm³ and platelets recover to a level >100,000 cells/mm³. In patients with NSCLC, resume treatment with every-3-week cycles at reduced doses for both weekly Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and every-3-week carboplatin after ANC recovers to at least 1500 cells/mm³ and platelet count of at least 100,000 cells/mm³ on Day 1 or to an ANC of at least 500 cells/mm³ and platelet count of at least 50,000 cells/mm³ on Days 8 or 15 of the cycle *[see Dosage and Administration (2.6)]*.

In patients with adenocarcinoma of the pancreas, withhold Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine if ANC is less than 1500 cells/mm³ or platelets are less than 50,000 cells/mm³ and delay initiation of the next cycle of therapy until ANC is at least 1500 cells/mm³ and platelet is less than 100,000 cells/mm³ on Day 1 of the cycle. Resume treatment with appropriate dose reduction if recommended *[see Dosage and Administration (2.6)]*.

5.2 Severe Neuropathy

Sensory neuropathy is dose- and schedule-dependent *[see Adverse Reactions (6.1)]*. If ≥ Grade 3 sensory neuropathy develops, withhold Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) treatment until resolution to Grade 1 or 2 for metastatic breast cancer or Grade 1 for NSCLC and pancreatic cancer followed by a dose reduction for all subsequent courses. Patients with NSCLC and pancreatic cancer followed by a dose reduction for Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Dosage and Administration (2.6)]*.

5.3 Sepsis

Sepsis occurred in 5% of patients with or without neutropenia who received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine. Biliary obstruction or presence of biliary stent were risk factors for severe or fatal sepsis.

If a patient becomes febrile (regardless of ANC) initiate treatment with broad spectrum antibiotics. For febrile neutropenia, interrupt Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine if ANC is less than 500 cells/mm³ and ANC ≥ 1500, then resume treatment at reduced dose levels *[see Dosage and Administration (2.6)]*.

5.4 Pneumonitis

Pneumonitis, including some cases that were fatal, occurred in 4% of patients receiving Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine.

Monitor patients for signs and symptoms of pneumonitis and interrupt Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine during evaluation of suspected pneumonitis. After ruling out infectious etiology and upon making a diagnosis of pneumonitis, permanently discontinue treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine.

5.5 Severe Hypersensitivity

Severe and sometimes fatal hypersensitivity reactions, including anaphylactic reactions, have been reported. Do not rechallenge patients who experience a severe hypersensitivity reaction to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) with this drug *[see Contraindications (4)]*.

Cross-hypersensitivity between Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and other taxane products has been reported and may include severe reactions such as anaphylaxis. Closely monitor patients with a previous history of hypersensitivity to other taxanes during initiation of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy.

5.6 Use in Patients with Hepatic Impairment

The exposure and toxicity of paclitaxel can be increased in patients with hepatic impairment. Closely monitor patients with mild hepatic impairment for severe myelosuppression.

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is not recommended for use in patients with AST > 10 x ULN; or bilirubin > 5 x ULN or with metastatic adenocarcinoma of the pancreas who have moderate to severe hepatic impairment. For MBC or NSCLC, reduce starting dose in patients with moderate to severe hepatic impairment *[see Dosage and Administration (2.5), Use in Specific Populations (8.7), Clinical Pharmacology (12.3)]*.

Albumin-bound particles for injectable suspension: white to yellow, sterile lyophilized powder containing 100 mg of paclitaxel formulated as albumin-bound particles in single-dose vial for reconstitution.

4 CONTRAINDICATIONS

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is contraindicated in patients with:

- Baseline neutrophil counts of < 1,500 cells/mm³ *[see Warnings and Precautions (5.1)]*.
- A history of severe hypersensitivity reactions to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Warnings and Precautions (5.5)]*.

5 WARNINGS AND PRECAUTIONS

5.1 Severe Myelosuppression

Severe myelosuppression (primarily neutropenia) is dose-dependent and is the dose-limiting toxicity of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). In clinical studies, Grade 4 neutropenia occurred in 34% of patients with metastatic breast cancer (MBC), 47% of patients with non-small cell lung cancer (NSCLC), and 38% of patients with pancreatic cancer.

Monitor for severe neutropenia and thrombocytopenia by performing complete blood cell counts frequently, including prior to dosing on Day 1 (for MBC) and Days 1, 8, and 15 (for NSCLC and for pancreatic cancer). Do not administer Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) to patients with baseline absolute neutrophil counts (ANC) of less than 1,500 cells/mm³ *[see Contraindications (4)]*.

In the case of severe neutropenia (<500 cells/mm³ for seven days or more) during a course of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy, reduce the dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in subsequent courses in patients with either MBC or NSCLC.

In patients with MBC, resume treatment with every-3-week cycles of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) after ANC recovers to at least 1,500 cells/mm³ and platelets recover to a level >100,000 cells/mm³. In patients with NSCLC, resume treatment with every-3-week cycles at reduced doses for both weekly Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and every-3-week carboplatin after ANC recovers to at least 1500 cells/mm³ and platelet count of at least 100,000 cells/mm³ on Day 1 or to an ANC of at least 500 cells/mm³ and platelet count of at least 50,000 cells/mm³ on Days 8 or 15 of the cycle *[see Dosage and Administration (2.6)]*.

In patients with adenocarcinoma of the pancreas, withhold Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine if ANC is less than 1500 cells/mm³ or platelets are less than 50,000 cells/mm³ and delay initiation of the next cycle of therapy until ANC is at least 1500 cells/mm³ and platelet is less than 100,000 cells/mm³ on Day 1 of the cycle. Resume treatment with appropriate dose reduction if recommended *[see Dosage and Administration (2.6)]*.

5.2 Severe Neuropathy

Sensory neuropathy is dose- and schedule-dependent *[see Adverse Reactions (6.1)]*. If ≥ Grade 3 sensory neuropathy develops, withhold Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) treatment until resolution to Grade 1 or 2 for metastatic breast cancer or Grade 1 for NSCLC and pancreatic cancer followed by a dose reduction for all subsequent courses. Patients with NSCLC and pancreatic cancer followed by a dose reduction for Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Dosage and Administration (2.6)]*.

5.3 Sepsis

Sepsis occurred in 5% of patients with or without neutropenia who received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine. Biliary obstruction or presence of biliary stent were risk factors for severe or fatal sepsis.

If a patient becomes febrile (regardless of ANC) initiate treatment with broad spectrum antibiotics. For febrile neutropenia, interrupt Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine if ANC is less than 500 cells/mm³ and ANC ≥ 1500, then resume treatment at reduced dose levels *[see Dosage and Administration (2.6)]*.

5.4 Pneumonitis

Pneumonitis, including some cases that were fatal, occurred in 4% of patients receiving Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with gemcitabine.

Monitor patients for signs and symptoms of pneumonitis and interrupt Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine during evaluation of suspected pneumonitis. After ruling out infectious etiology and upon making a diagnosis of pneumonitis, permanently discontinue treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine.

5.5 Severe Hypersensitivity

Severe and sometimes fatal hypersensitivity reactions, including anaphylactic reactions, have been reported. Do not rechallenge patients who experience a severe hypersensitivity reaction to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) with this drug *[see Contraindications (4)]*.

Cross-hypersensitivity between Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and other taxane products has been reported and may include severe reactions such as anaphylaxis. Closely monitor patients with a previous history of hypersensitivity to other taxanes during initiation of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) therapy.

5.6 Use in Patients with Hepatic Impairment

The exposure and toxicity of paclitaxel can be increased in patients with hepatic impairment. Closely monitor patients with mild hepatic impairment for severe myelosuppression.

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is not recommended for use in patients with AST > 10 x ULN or with metastatic adenocarcinoma of the pancreas who have moderate to severe hepatic impairment. For MBC or NSCLC, reduce starting dose in patients with moderate to severe hepatic impairment *[see Dosage and Administration (2.5), Use in Specific Populations (8.7), Clinical Pharmacology (12.3)]*.

Table 2: Dose Recommendation and Modifications for Neutropenia and/or Thrombocytopenia at the Start of a Cycle or within a Cycle for Patients with Adenocarcinoma of the Pancreas

Cycle Day	ANC (cells/mm³)		Platelet count (cells/mm³)	Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) / Gemcitabine
Day 15: If Day 8 doses were reduced or given without modification:				
	500 to < 1000	OR	50,000 to < 75,000	Reduce 1 dose level from Day 1
	< 500	OR	< 50,000	Withhold doses
Day 15: If Day 8 doses were withheld:				
	≥ 1000	OR	≥ 75,000	Reduce 1 dose level from Day 1
	500 to < 1000	OR	50,000 to < 75,000	Reduce 2 dose levels from Day 1
	< 500	OR	< 50,000	Withhold doses

ANC = Absolute Neutrophil Count

Recommended dose modifications for other adverse reactions in patients with pancreatic cancer are provided in Table 5.

Table 5: Dose Modifications for Other Adverse Reactions in Patients with Adenocarcinoma of the Pancreas

Adverse Reaction	Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)	Gemcitabine
Febrile Neutropenia: Grade 3 or 4	Withhold until fever resolves and ANC ≥ 1500; resume at next lower dose level	
Peripheral Neuropathy: Grade 3 or 4	Withhold until improves to ≤ Grade 1; resume at next lower dose level	No dose reduction
Cutaneous Toxicity: Grade 2 or 3	Reduce to next lower dose level; discontinue treatment if toxicity persists	
Gastrointestinal Toxicity: Grade 3 mucositis or diarrhea	Withhold until improves to ≤ Grade 1; resume at next lower dose level	

2.7 Preparation for Intravenous Administration

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is a cytotoxic drug. Follow applicable special handling and disposal procedures. The use of gloves is recommended. If Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) (lyophilized cake or reconstituted suspension) contacts the skin, wash the skin immediately and thoroughly with soap and water. Following topical exposure to paclitaxel, events may include tingling, burning, redness, and swelling. If Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) contacts mucous membranes, rinse thoroughly with water.

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is supplied as a sterile lyophilized powder for reconstitution before use.

Read the entire preparation instructions prior to reconstitution.

1. Aseptically, reconstitute each vial by injecting 20 mL of 0.9% Sodium Chloride Injection, USP.
2. Slowly inject the 20 mL of

(approximately 2% of the daily maximum recommended human dose on a mg/m² basis) caused embryo-fetal toxicities, as indicated by intrauterine mortality, increased resorptions (up to 5-fold), reduced numbers of litters and live fetuses, reduction in fetal body weight, and increase in fetal anomalies. Fetal anomalies included soft tissue and skeletal malformations, such as eye bulge, folded retina, microphthalmia, and dilation of brain ventricles.

8.2 Lactation

Risk Summary

There are no data on the presence of paclitaxel in human milk, or its effect on the breastfed child or on milk production. In animal studies, paclitaxel and/or its metabolites were excreted into the milk of lactating rats (see Data). Because of the potential for serious adverse reactions in a breastfed child from Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound), advise lactating women not to breastfeed during treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and for two weeks after the last dose.

Data

Animal Data

Following intravenous administration of radiolabeled paclitaxel to rats on days 9 to 10 postpartum, concentrations of radioactivity in milk were higher than in plasma and declined in parallel with the plasma concentrations.

8.3 Females and Males of Reproductive Potential

Based on animal studies and mechanism of action, Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) can cause fetal harm when administered to a pregnant woman *[see Use in Specific Populations (8.1)]*.

Pregnancy Testing

Verify the pregnancy status of females of reproductive potential prior to starting treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound).

Contraception

Females

Advise females of reproductive potential to use effective contraception and avoid becoming pregnant during treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and for at least six months after the last dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound).

Males

Based on findings in genetic toxicity and animal reproduction studies, advise males with female partners of reproductive potential to use effective contraception and avoid fathering a child during treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and for at least three months after the last dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *(see Use in Specific Populations (8.1) and Nonclinical Toxicology (13.1))*.

Infertility

Females and Males

Based on findings in animals, Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) may impair fertility in females and males of reproductive potential *[see Nonclinical Toxicology (13.1)]*.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established. Pharmacokinetics, safety, and antitumor activity of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) were assessed in an open-label, dose escalation, dose expansion study (NCT01992103) in 96 pediatric patients aged 1.4 to < 17 years with recurrent or refractory pediatric solid tumors. The maximum tolerated dose (MTD) normalized for body surface area (BSA) was lower in pediatric patients compared to adults. No new safety signals were observed in pediatric patients across these studies.

Paclitaxel protein-bound exposures normalized by dose were higher in 96 pediatric patients (aged 1.4 to < 17 years) as compared to those in adults.

8.5 Geriatric Use

Of the 229 patients in the randomized study who received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) for the treatment of metastatic breast cancer, 13% were at least 65 years of age and < 2% were 75 years or older. This study of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) did not include a sufficient number of patients with metastatic breast cancer who were 65 years and older to determine whether they respond differently from younger patients.

A subsequent pooled analysis was conducted in 981 patients receiving Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) monotherapy for metastatic breast cancer, of which 15% were 65 years of age or older and 2% were 75 years of age or older. A higher incidence of epistaxis, diarrhea, dehydration, fatigue, and peripheral edema was found in patients 65 years of age or older.

Of the 514 patients in the randomized study who received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and carboplatin for the first-line treatment of non-small cell lung cancer, 31% were 65 years or older and 3.5% were 75 years or older. Myelosuppression, peripheral neuropathy, and arthralgia were more frequent in patients 65 years or older compared to patients younger than 65 years old. No overall difference in effectiveness, as measured by response rates, was observed between patients 65 years or older compared to patients younger than 65 years old.

Of the 431 patients in the randomized study who received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and gemcitabine for the first-line treatment of pancreatic adenocarcinoma, 41% were 65 years or older and 10% were 75 years or older. No overall differences in effectiveness were observed between patients who were 65 years of age or older and younger patients. Diarrhea, decreased appetite, dehydration, and epistaxis were more frequent in patients 65 years or older compared with patients younger than 65 years old. Clinical studies of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) did not include sufficient number of patients with pancreatic cancer who were 75 years and older to determine whether they respond differently from younger patients.

8.6 Renal Impairment

No adjustment of the starting Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) dose is required for patients with mild to moderate renal impairment (estimated creatinine clearance 30 to <90 mL/min) *[see Clinical Pharmacology (12.3)]*. There are insufficient data to permit dosage recommendations in patients with severe renal impairment or end stage renal disease (estimated creatinine clearance <30 mL/min).

8.7 Hepatic Impairment

No adjustment of the starting Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) dose is required for patients with mild hepatic impairment (total bilirubin > ULN and ≤ 1.5 x ULN and aspartate aminotransferase [AST] ≤ 10 x ULN). Reduce Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) starting dose in patients with moderate to severe hepatic impairment *[see Dosage and Administration (2.5) and Clinical Pharmacology (12.3)]*. Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is not recommended for use in patients with total bilirubin > 5 x ULN or AST > 10 x ULN *[see Dosage and Administration (2.5), Warnings and Precautions (5.6), and Clinical Pharmacology (12.3)]*. Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is not recommended for use in patients with metastatic adenocarcinoma of the pancreas who have moderate to severe hepatic impairment *[see Dosage and Administration (2.5)]*.

10 OVERDOSAGE

There is no known antidote for Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) overdosage. The primary anticipated complications of overdosage would consist of bone marrow suppression, sensory neurotoxicity, and mucositis.

11 DESCRIPTION

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is paclitaxel formulated as albumin-bound nanoparticles with a mean particle size of approximately 130 nanometers. Paclitaxel exists in the particles in a non-crystalline, amorphous state. Paclitaxel is a microtubule inhibitor. The chemical name for paclitaxel is 5β,20-Epoxy-1,2α,4,7β,10,13α-hexahydroxytax-11-en-9-one 4,10-diacetate 2-benzoate 13-ester with (2R,3S)-N-benzoyl-3-phenylisoserine. The empirical formula is $C_{47}H_{51}NO_{14}$, and the molecular weight is 853.91. Paclitaxel has the following structural formula:



Paclitaxel is a white to off-white crystalline powder. It is highly lipophilic, insoluble in water, and melts at approximately 216°C to 217°C.

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is supplied as a white to yellow, sterile, lyophilized powder for reconstitution with 20 mL of 0.9% Sodium Chloride Injection, USP prior to intravenous infusion. Each single-dose vial contains 100 mg of paclitaxel (bound to human albumin) and approximately 900 mg of human albumin (containing sodium caprylate and sodium acetyltryptophanate). Each milliliter (mL) of reconstituted suspension contains 5 mg paclitaxel formulated as albumin-bound particles. Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is free of solvents.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is a microtubule inhibitor that promotes the assembly of microtubules from tubulin dimers and stabilizes microtubules by preventing depolymerization. This stability results in the inhibition of the normal dynamic reorganization of the microtubule network that is essential for vital interphase and mitotic cellular functions. Paclitaxel induces abnormal arrays or "bundles" of microtubules throughout the cell cycle and multiple asters of microtubules during mitosis.

12.3 Pharmacokinetics

The pharmacokinetics of total paclitaxel following 30- and 180-minute infusions of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) dose levels of 80 to 375 mg/m² (0.31 to 1.15 times the maximum approved recommended dosage) were determined in clinical studies. Dose levels of mg/m² refer to mg of paclitaxel in Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound). Following intravenous administration of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) to patients with solid tumors, paclitaxel plasma concentrations declined in a biphasic manner, the initial rapid decline representing distribution to the peripheral compartment and the slower second phase representing drug elimination.

Following Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) infusion, paclitaxel exhibited linear drug exposure (AUC) across clinical doses ranging from 80 to 300 mg/m² (0.31 to 1.15 times the maximum approved recommended dosage). The pharmacokinetics of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) were independent of the duration of intravenous administration.

The pharmacokinetic data of 260 mg/m² Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) administered over a 30-minute infusion was compared to the pharmacokinetics of 175 mg/m² paclitaxel injection over a 3-hour infusion. Clearance was larger (43%) and the volume of distribution was higher (53%) for Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) than for paclitaxel injection. There were no differences in terminal half-lives.

Distribution

Following Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) administration to patients with solid tumors, paclitaxel is evenly distributed into blood cells and plasma and is highly bound to plasma proteins (94%). The total volume of distribution is approximately 1741 L, the large volume of distribution indicates extensive extravascular distribution and/or tissue binding of paclitaxel.

In a within-patient comparison study, the fraction of unbound paclitaxel in plasma was significantly higher with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) (6.2%) than with solvent-based paclitaxel (2.3%). This contributes to significantly higher exposure to unbound Paclitaxel with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) compared with solvent-based paclitaxel, when the total exposure is comparable. *In vitro* studies of binding to human serum proteins, using paclitaxel concentrations ranging from 0.1 to 50 μg/mL, indicated that the presence of cimetidine, ranitidine, dexamethasone, or diphenhydramine did not affect protein binding of paclitaxel.

Elimination

At the clinical dose range of 80 to 300 mg/m² (0.31 to 1.15 times the maximum approved recommended dosage), the mean total clearance of paclitaxel ranges from 13 to 30 L/h/m² and the mean terminal half-life ranges from 13 to 27 hours.

Metabolism

In vitro studies with human liver microsomes and tissue slices showed that paclitaxel in Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) was metabolized primarily to 6α-hydroxypaclitaxel by CYP2C8; and to two minor metabolites, 3′-p-hydroxypaclitaxel and 6α, 3′-p-dihydroxypaclitaxel, by CYP3A4. *In vitro*, the metabolism of paclitaxel to 6α-hydroxypaclitaxel was inhibited by a number of agents (ketoconazole, verapamil, diazepam, quinidine, dexamethasone, cyclosporin, teniposide, etoposide, and vincristine), but the concentrations used exceeded those found *in vivo* following normal therapeutic doses. Testosterone, 17α-ethinyl estradiol, retinoic acid, and quercetin, a specific inhibitor of CYP2C8, also inhibited the formation of 6α-hydroxypaclitaxel *in vitro*. The pharmacokinetics of paclitaxel may also be altered *in vivo* as a result of interactions with compounds that are substrates, inducers, or inhibitors of CYP2C8 and/or CYP3A4 *[see Drug Interactions (7)]*.

Excretion

After a 30-minute infusion of 260 mg/m² doses of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound), the mean values for cumulative urinary recovery of unchanged drug (4%) indicated extensive non-renal clearance. Less than 1% of the total administered dose was excreted in urine as the metabolites 6α-hydroxypaclitaxel and 3′-p-hydroxypaclitaxel. Fecal excretion was approximately 20% of the total dose administered.

Specific Populations

No clinically meaningful differences in the pharmacokinetics of paclitaxel in Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) were observed based on body weight (40 to 143 kg), body surface area (1.3 to 2.4 m²), sex, race (Asian vs. White), age (24 to 85 years), type of solid tumors, mild to moderate renal impairment (creatinine clearance 30 to <90 mL/min), and mild hepatic impairment (total bilirubin >1 to ≤1.5 x ULN and AST ≤10 x ULN).

Patients with moderate (total bilirubin >1.5 to 3 x ULN and AST ≤10 x ULN) or severe (total bilirubin >3 to 5 x ULN) hepatic impairment had a 22% to 26% decrease in the maximum elimination rate of paclitaxel and approximately 20% increase in mean paclitaxel AUC compared with patients with normal hepatic function (total bilirubin ≤ULN and AST ≤ULN) *[see Dosage and Administration (2.5) and Use in Specific Populations (8.7)]*.

The effect of severe renal impairment or end stage renal disease (creatinine clearance < 30 mL/min) on the pharmacokinetics of paclitaxel in Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is unknown.

Drug Interaction Studies

Carboplatin: Administration of carboplatin immediately after the completion of the Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) infusion to patients with NSCLC did not cause clinically meaningful changes in paclitaxel exposure. The observed mean AUC_{0-t} of free carboplatin was approximately 23% higher than the targeted value (6 min*mg/mL), but its mean half-life and clearance were consistent with those reported in the absence of paclitaxel.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) has not been studied.

Paclitaxel was clastogenic *in vitro* (chromosome aberrations in human lymphocytes) and *in vivo* (micronucleus test in mice). Paclitaxel was not mutagenic in the Ames test or the CHO/HGPRT gene mutation assay.

Administration of paclitaxel formulated as albumin-bound particles to male rats at 42 mg/m² on a weekly basis (approximately 16% of the daily maximum recommended human exposure on a body surface area basis) for 11 weeks prior to mating with untreated female rats resulted in significantly reduced fertility accompanied by decreased pregnancy rates and increased loss of embryos in mated females. A dose of 42 mg/m² also reduced male reproductive organ weights, mating performance, and sperm production. Testicular atrophy/degeneration was observed in single-dose toxicology studies in animals administered paclitaxel formulated as albumin-bound particles at doses lower than the recommended human dose; doses were 54 mg/m² in rodents and 175 mg/m² in dogs. Similar testicular degeneration was seen in monkeys administered three weekly doses of 108 mg/m² paclitaxel formulated as albumin bound particles.

Administration of paclitaxel prior to and during mating produced impairment of fertility in male and female rats. Paclitaxel caused reduced fertility and reproductive indices, and increased embryo-fetal toxicity.

14 CLINICAL STUDIES

14.1 Metastatic Breast Cancer

Data from 106 patients accrued in two single arm open label studies and from 460 patients enrolled in a randomized comparative study were available to support the use of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in metastatic breast cancer.

Single Arm Open Label Studies

In one study, Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) was administered as a 30-minute infusion at a dose of 175 mg/m² to 43 patients with metastatic breast cancer. The second trial utilized a dose of 300 mg/m² as a 30-minute infusion in 63 patients with metastatic breast cancer. Cycles were administered at 3-week intervals. Objective responses were observed in both studies.

Randomized Comparative Study

This multicenter trial was conducted in 460 patients with metastatic breast cancer. Patients were randomized to receive Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) at a dose of 260 mg/m² given as a 30-minute infusion, or paclitaxel injection at 175 mg/m² given as a 3-hour infusion. Sixty-four percent of patients had impaired performance status (ECOG 1 or 2) at study entry; 79% had visceral metastases; and 76% had > 3 sites of metastases. Fourteen percent of the patients had not received prior chemotherapy; 27% had received chemotherapy in the adjuvant setting, 40% in the metastatic setting and 19% in both metastatic and adjuvant settings. Fifty-nine percent received study drug as second or greater than second-line therapy. Seventy-seven percent of the patients had been previously exposed to anthracyclines.

In this trial, patients in the Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) treatment arm had a statistically significantly higher reconciled target lesion response rate (the trial primary endpoint) of 21.5% (95% CI: 16.2% to 26.7%), compared to 11.1% (95% CI: 6.9% to 15.1%) for patients in the paclitaxel injection treatment arm. See Table 11. There was no statistically significant difference in overall survival between the two study arms.

Table 11: Efficacy Results from Randomized Metastatic Breast Cancer Trial

	Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) 260 mg/m²	Paclitaxel Injection 175 mg/m²
Reconciled Target Lesion Response Rate (primary endpoint)[a]		
All randomized patients	Response Rate [95% CI]	50/233 (21.5%) [16.19% – 26.73%]	25/227 (11.1%) [6.94% – 15.09%]
	p-value[b]	0.003	
Patients who had failed combination chemotherapy or relapsed within 6 months of adjuvant chemotherapy[c]	Response Rate [95% CI]	20/129 (15.5%) [9.26% – 21.75%]	12/143 (8.4%) [3.85% – 12.94%]

[a] Reconciled Target Lesion Response Rate (TLRR) was the prospectively defined protocol specific endpoint, based on independent radiologic assessment of tumor responses reconciled with investigator responses (which also included clinical information) for the first 6 cycles of therapy. The reconciled TLRR was lower than the investigator Reported Response Rates, which are based on all cycles of therapy.
[b] From Cochran-Mantel-Haenszel test stratified by 1st line vs. > 1st line therapy.
[c] Prior therapy included an anthracycline unless clinically contraindicated.

14.2 Non-Small Cell Lung Cancer

A multicenter, randomized, open-label study was conducted in 1052 chemotherapy naive patients with Stage IIIb/IV non-small cell lung cancer to compare Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in combination with carboplatin to paclitaxel injection in combination with carboplatin as first-line treatment in patients with advanced non-small cell lung cancer. Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) was administered as an intravenous infusion over 30 minutes at a dose of 100 mg/m² on Days 1, 8, and 15 of each 21-day cycle. Paclitaxel injection was administered as an intravenous infusion over 3 hours at a dose of 200 mg/m², following premedication. In both treatment arms carboplatin at a dose of AUC = 6 mg•min/mL was administered intravenously on Day 1 of each 21-day cycle after completion of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)/paclitaxel infusion. Treatment was administered until disease progression or development of an unacceptable toxicity. The major efficacy outcome measure was overall response rate as determined by a central independent review committee using RECIST guidelines (Version 1.0).

In the intent-to-treat (all-randomized) population, the median age was 60 years; 75% were men, 81% were White, 49% had adenocarcinoma, 43% had squamous cell lung cancer, 76% were ECOG PS 1, and 73% were current or former smokers. Patients received a median of 6 cycles of treatment in both study arms.

Patients in the Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)/carboplatin arm had a statistically significantly higher overall response rate compared to patients in the paclitaxel injection/carboplatin arm [(33% versus 25%) see Table 12]. There was no statistically significant difference in overall survival between the two study arms.

Table 12: Efficacy Results from Randomized Non-Small Cell Lung Cancer Trial (Intent-to-Treat Population)

	Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) (100 mg/m² weekly) + carboplatin (N=521)	Paclitaxel Injection (200 mg/m² every 3 weeks) + carboplatin (N=531)
Overall Response Rate (ORR)		
Confirmed complete or partial overall response, n (%)	170 (33%)	132 (25%)
95% CI	28.6, 36.7	21.2, 28.5
P-value (Chi-Square test)	0.005	
Median DoR in months (95% CI)	6.9 (5.6, 8.0)	6.0 (5.6, 7.1)
Overall Response Rate by Histology		
Carcinoma/ Adenocarcinoma	66/254 (26%)	71/264 (27%)
Squamous Cell Carcinoma	94/229 (41%)	54/221 (24%)
Large Cell Carcinoma	3/9 (33%)	2/13 (15%)
Other	7/29 (24%)	5/33 (15%)

CI = confidence interval; DoR= Duration of response.

14.3 Adenocarcinoma of the Pancreas

A multicenter, multinational, randomized, open-label study was conducted in 861 patients comparing Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) plus gemcitabine versus gemcitabine monotherapy as first-line treatment of metastatic adenocarcinoma of the pancreas. Key eligibility criteria were Karnofsky Performance Status (KPS) ≥ 70, normal bilirubin level, transaminase levels ≤ 2.5 times the upper limit of normal (ULN) or ≤ 5 times the ULN for patients with liver metastasis, no prior cytotoxic chemotherapy in the adjuvant setting or for metastatic disease, no ongoing active infection requiring systemic therapy, and no history of interstitial lung disease. Patients with rapid decline in KPS (≥ 10%) or serum albumin (≥ 20%) during the 14 day screening period prior to study randomization were ineligible.

A total of 861 patients were randomized (1:1) to the Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)/gemcitabine arm (N=431) or to the gemcitabine arm (N=430).

Randomization was stratified by geographic region (Australia, Western Europe, Eastern Europe, or North America), KPS (70 to 80 versus 90 to 100), and presence of liver metastasis (yes versus no). Patients randomized to Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)/gemcitabine received Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) 125 mg/m² as an intravenous infusion over 30-40 minutes followed by gemcitabine 1000 mg/m² as an intravenous infusion over 30-40 minutes on Days 1, 8, and 15 of each 28-day cycle. Patients randomized to gemcitabine received 1000 mg/m² as an intravenous infusion over 30-40 minutes weekly for 7 weeks followed by a 1-week rest period in Cycle 1 then as 1000 mg/m² on Days 1, 8, and 15 of each subsequent 28-day cycle. Patients in both arms received treatment until disease progression or unacceptable toxicity. The major efficacy outcome measure was overall survival (OS). Additional outcome measures were progression-free survival (PFS) and overall response rate (ORR), both assessed by independent, central, blinded radiological review using RECIST (version 1.0).

In the intent-to-treat (all randomized) population, the median age was 63 years (range 27-88 years) with 42% ≥ 65 years of age; 58% were men; 93% were White and the KPS was 90-100 in 60%. Disease characteristics included 46% of patients with 3 or more metastatic sites; 84% of patients had liver metastasis; and the location of the primary pancreatic lesion was in the head of pancreas (43%), body (31%), or tail (25%).

Results for overall survival, progression-free survival, and overall response rate are shown in Table 13.

Table 13: Efficacy Results from Randomized Study in Patients with Adenocarcinoma of the Pancreas (ITT Population)

	Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) (125 mg/m²) and gemcitabine (N = 431)	Gemcitabine (N = 430)
Overall Survival		
Number of deaths, n (%)	333 (77)	359 (83)
Median Overall Survival (months)	**8.5**	**6.7**
95% CI	7.9, 9.5	6.0, 7.2
HR (95% CI) [a]	0.72 (0.62, 0.83)	
P-value[b]	<0.0001	
Progression-free Survival[c]		
Death or progression, n (%)	277 (64)	265 (62)
Median Progression-free Survival (months)	**5.5**	**3.7**
95% CI	4.5, 5.9	3.6, 4.0
HR (95% CI) [a]	0.69 (0.58, 0.82)	
P-value[b]	<0.0001	
Overall Response Rate[c]		
Confirmed complete or partial overall response, n (%)	**99 (23)**	**31 (7)**
95% CI	19.1, 27.2	5.0, 10.1
P-value [d]	<0.0001	

CI = confidence interval; HR = hazard ratio of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) plus gemcitabine / gemcitabine, ITT = intent-to-treat population.
[a] Stratified Cox proportional hazard model.
[b] Stratified log-rank test stratified by geographic region (North America versus Others), Karnofsky performance score (70 to 80 versus 90 to 100), and presence of liver metastasis (yes versus no).
[c] Based on Independent Radiological Reviewer Assessment.
[d] Chi-square test.

In exploratory analyses conducted in clinically relevant subgroups with a sufficient number of subjects, the treatment effects on overall survival were similar to that observed in the overall study population.

Figure 1: Kaplan-Meier Curve of Overall Survival(Intent-to-TreatPopulation)



15 REFERENCES

1. OSHA Hazardous Drugs. OSHA http://www.osha.gov/SLTC/hazardousdrugs/index.html

16 HOW SUPPLIED/STORAGE AND HANDLING

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is a white to to yellow, sterile lyophilized powder supplied as:

NDC 24979-710-51 100 mg of paclitaxel in a single-dose vial, individually packaged in a carton.

Store the vials in original cartons at 20°C to 25°C (68°F to 77°F). Retain in the original package to protect from bright light.

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) is a cytotoxic drug. Follow applicable special handling and disposal procedures.[1]

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the approved patient labeling (Patient Information).

Severe Myelosuppression

- Patients must be informed of the risk of low blood cell counts and severe and life-threatening infections and instructed to contact their healthcare provider immediately for fever or evidence of infection *[see Warnings and Precautions (5.1), (5.3)]*.

Severe Neuropathy

- Patients must be informed that sensory neuropathy occurs frequently with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and patients should advise their healthcare providers of numbness, tingling, pain, or weakness involving the extremities *[see Warnings and Precautions (5.2)]*.

Pneumonitis

- Instruct patients to contact their healthcare provider immediately for sudden onset of dry persistent cough, or shortness of breath *[see Warnings and Precautions (5.4)]*.

Severe Hypersensitivity

- Instruct patients to contact their healthcare provider for signs of an allergic reaction, which could be severe and sometimes fatal *[see Warnings and Precautions (5.5)]*.

Common Adverse Reactions

- Explain to patients that alopecia, fatigue/asthenia, and myalgia/arthralgia occur frequently with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound).
- Instruct patients to contact their healthcare providers for persistent vomiting, diarrhea, or signs of dehydration *[see Adverse Reactions (6)]*.

Embryo-Fetal Toxicity

- Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) injection can cause fetal harm. Advise patients to avoid becoming pregnant while receiving this drug. Females of reproductive potential should use effective contraception during treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and for at least six months after the last dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Warnings and Precautions (5.8) and Use in Specific Populations (8.1, 8.3)]*.
- Advise male patients with female partners of reproductive potential to use effective contraception and avoid fathering a child during treatment with Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and for at least three months after the last dose of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) *[see Use in Specific Populations (8.3)]*.

Lactation

- Advise patients not to breastfeed while taking Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) and for two weeks after receiving the last dose *[see Use in Specific Populations (8.2)]*.

Infertility

- Advise males and females of reproductive potential that Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) may impair fertility *[see Use in Specific Populations (8.3)]*.

Manufactured for: TWi Pharmaceuticals USA, Inc.
Paramus, NJ 07652

PACTWIPI.012/PPI.012

Manufactured for:
TWi Pharmaceuticals USA, Inc.
Paramus, NJ 07652

PACTWIPPI.012

For more information, call 1-844-518-2989

Distributed by
The U.S. Food and Drug Administration

Revised: 8/2020

The most common side effects of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in people with breast cancer include:

- hair loss
- numbness, tingling, pain or weakness in your hands or feet
- tiredness
- changes in your liver function tests
- nausea
- infections
- decreased white blood cell count
- abnormal heartbeat
- decreased red blood cell count (anemia)
- low red blood cell count (anemia) may cause you to feel weak, tired, or short of breath

The most common side effects of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in people with non-small cell lung cancer include:

- low red blood cell count (anemia)
- numbness, tingling pain, or weakness in your hands or feet
- tiredness
- decreased white blood cell count
- nausea

The most common side effects of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in people with pancreatic cancer include:

- decreased white blood cell count
- tiredness, weakness, or lack of energy
- nausea
- vomiting
- hair loss
- diarrhea
- rash
- tiredness
- fever
- decreased appetite
- signs of dehydration including decreased urination, headache, or muscle cramps

Tell your healthcare provider if you have vomiting, diarrhea, or signs of dehydration that does not go away.

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) may cause birth defects. Advise females and males with partners who may become pregnant that this may affect your ability to have a child. Talk to your healthcare provider if this is a concern for you. Please read all of the information about Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) that is written for health professionals.

What are the ingredients in Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)?

Active ingredient: paclitaxel (bound to human albumin).

Other ingredient: human albumin (containing sodium caprylate and sodium acetyltryptophanate).

General information about the safe and effective use of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound).

Medicines are sometimes prescribed for conditions that are not mentioned in a Patient Information leaflet. You can ask your healthcare provider or pharmacist for information about Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) that is written for health professionals.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

The most common side effects of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in people with pancreatic cancer include:

- decreased white blood cell count
- tiredness, weakness, or lack of energy
- nausea
- vomiting
- hair loss
- diarrhea
- rash
- tiredness
- fever
- decreased appetite
- signs of dehydration including decreased urination, headache, or muscle cramps

Tell your healthcare provider if you have vomiting, diarrhea, or signs of dehydration that does not go away.

Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) may cause birth defects. Advise females and males with partners who may become pregnant that this may affect your ability to have a child. Talk to your healthcare provider if this is a concern for you. Please read all of the information about Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) that is written for health professionals.

What are the ingredients in Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound)?

Active ingredient: paclitaxel (bound to human albumin).

Other ingredient: human albumin (containing sodium caprylate and sodium acetyltryptophanate).

General information about the safe and effective use of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound).

Medicines are sometimes prescribed for conditions that are not mentioned in a Patient Information leaflet. You can ask your healthcare provider or pharmacist for information about Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) that is written for health professionals.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

The most common side effects of Paclitaxel Protein-Bound Particles for Injectable Suspension (Albumin-Bound) in people with breast cancer include:

- hair loss
- numbness, tingling, pain or weakness in your hands or feet
- tiredness
- changes in your liver function tests
- nausea
- infections
- decreased white blood cell count
- abnormal heartbeat
- decreased red blood cell count (anemia)
- low red blood cell count (anemia) may cause you to feel weak, tired, or short of breath

Nosco
complete packaging | individual solutions™

NOSCO GRAPHIC SERVICES
11200 88th Avenue, Pleasant Prairie, WI 53158 | Phone: 847.336.4200
Serving Nosco Facilities in Illinois, Texas, Pennsylvania and New York.

Proof A
12/20/2021

Product Information	Specified Colors	Approved By
Item: 112462 **Eval:** X5122547 Cust: CELGENE Size: +39. x +20.5	Black Head to Head Item Group: 5105x90	Name Date ☐ OK to Print ☐ New Proof Required We must have signed proof before we can begin production. This proof is to show size, copy placement and color break. Spot PMS colors will be matched on press to PMS Book, 4 Color Process Digital Color Guide, Extended Gamut Digital Color Guide and/or approved color standards. 4 Color Process proofs are verified to G7 standards and approved for color match.

PDF/X overprint ☒ ☐ ☒ ☐ When correctly rendered in compliance with PDF/X standards no 'x' should appear in any box. Please check overprint settings should you see an 'x'.